Assessing the Antioxidant Activity of Dexmedetomidine by Measuring the Plasma Antioxidant Enzyme Activity of Catalase, Glutathione Peroxidase, and Superoxide Dismutase During Lumbar Spine Laminectomy

Kamran Mottaghi ¹, Masoud Nashibi ², Sogol Asgari ², Farahd Safari ², Matin Bahrebar ¹ and Parisa Sezari ^{2,*}

¹Anesthesiology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Shahid Beheshti University of Medical Sciences, Tehran, Iran

* Corresponding author: Anesthesiology Department, Lohman Hakim Hospital, Shahid Beheshti University of Medical Sciences, 13336-35445, Tehran, Iran. Email: psezari@sbmu.ac.ir

Received 2021 July 25; Revised 2021 September 04; Accepted 2021 October 22.

Abstract

Background: Current knowledge on antioxidant properties of anesthetics is inconclusive. Nevertheless, experiments on different anesthesia-related drugs and techniques are growing.

Objectives: The aim of this prospective blinded randomized study was to assess the effect of dexmedetomidine (Dex) infusion during anesthesia on oxidative stress resulting from lumbar laminectomy.

Methods: The antioxidant activity of Dex compared to placebo (normal saline) was evaluated by measuring the antioxidant enzyme activity of catalase (CAT), superoxide dismutase (SOD), and glutathione peroxidase (GPX). A total of 56 patients who underwent laminectomy surgery were allocated into two groups of intervention and control. Dex group received $0.6 \mu\text{g.kg}^{-1}$ dexmedetomidine infusion before induction of anesthesia over 15 minutes, followed by $0.4 \mu\text{g.kg}^{-1}.\text{hr}^{-1}$ maintenance. Control group received the same volume and sequence of normal saline. Anesthesia protocol was similar in both groups. The antioxidant activity of the previously mentioned enzymes was measured at the beginning of the anesthesia (T1) and after surgery (T2). Propofol consumption, hemodynamic indices, shivering, bleeding volume, and pain numeric rating scale (NRS) were recorded.

Results: Although the serum levels of CAT and GPX increased during the procedure, these changes were not significant (P-values 0.579 and 0.762, respectively). Also, the mean SOD level did not change over time in any of the groups, and it was not meaningfully different between the groups at any of the predetermined times (P-value 0.665).

Conclusions: As expected, Dex decreased propofol consumption, mean heart rate, mean arterial pressure, and pain intensity all over the perioperative period. It seems that adding Dex to anesthesia regimen during laminectomy had no dominant antioxidant effects.

Keywords: Glutathione Peroxidase, Superoxide Dismutase, Catalase, Dexmedetomidine, Anesthesia, Laminectomy, Stress Response, Antioxidants

1. Background

Surgical procedures induce a stress response (1), which increase reactive oxygen species (ROS) production and promote neurodegeneration (2). Endogenous antioxidants such as glutathione peroxidase (GPX), superoxide dismutase (SOD), and catalase (CAT) are meant to counteract ROS (3).

Dexmedetomidine (DEX), an α_2 adrenoreceptor agonist with anxiolytic (4), sympatholytic (5, 6), cognitive advantages (7) and pulmonary-mechanics modifying effects (8), has been used to attenuate hemodynamic response

during anesthesia (9) and surgery. It has also been used to alleviate stress response and ROS surge.

Spinal surgeries are of moderate stress level, which may propagate ROS (10). Thus, different techniques and medications might be used to attenuate oxidative response.

2. Methods

2.1. Anesthesia Method

In this randomized, blinded clinical trial, 56 patients (age range: 18 - 65 years) were included according to

the classification of American Society of Anesthesiologists. The participants had no past medical history of spine surgery, body mass index (BMI) less than 30 kg.m⁻², no addiction to opioids, alcohol, narcotics, or any illicit drugs, and they were candidates for single level lumbar spine laminectomy. Exclusion criteria were bleeding more than 10% of patient's blood volume, duration of surgery more than four hours, change of surgical approach by the surgeon, and transfusion and inotrope administration during surgery. Four patients were excluded due to changes in surgical approach (decided by the surgeon during the procedure), and one patient was excluded following bleeding more than acceptable volume. Finally, 51 patients were included in the study.

Patients were allocated into two groups of Dex (D group; n = 25) and control (C group; n = 26). The surgery team and anesthesia technique were similar in both groups. Pulse oximetry (SpO₂), non-invasive blood pressure monitoring (NIBP), capnometry (ETCO₂), electrocardiogram (ECG), cerebral state monitoring (CSM), and train of four (TOF) were performed. Normal saline (5 mL/kg) was infused followed by midazolam (0.03 mg/kg) and fentanyl (3 mcg/kg) as premedication. General anesthesia was induced by injecting intravenous propofol (2 mg.kg⁻¹), cisatracurium (0.15 mg.kg⁻¹), and lidocaine (1 mg.kg⁻¹). Moreover, propofol (100 - 300 mcg.kg⁻¹.hr⁻¹) was used to maintain cerebral state index (CSI) between 40 and 60.

Dex group received dexmedetomidine (0.6 µg.kg⁻¹) as loading dose 15 minutes before induction of anesthesia, followed by 0.4 µg.kg⁻¹.hr⁻¹ as maintenance. Control group received the same volume and sequence of normal saline. Trachea was intubated using properly sized endotracheal armored tube. During surgery, 50 µg of fentanyl was injected every 45 minutes until the end of surgery. If heart rate dropped to less than 40 beat per minute, atropine 0.5 mg was injected intravenously, and if mean arterial pressure was less than 60 mmHg, 10 mg of ephedrine was administered. In case of refractory hypotension, Dex infusion was decreased by 0.1 mcg.kg⁻¹.hr⁻¹, further followed by 20 mcg.kg⁻¹.hr⁻¹ decrease in propofol infusion rate. In case of rising mean pressure to more than 20% of baseline measures, trinitroglycerin was administered (3 - 5 mcg.min⁻¹). Anesthesiologist and surgeon were not informed about the groups and were blinded to the study. Dex discontinued after the skin closure at the end of surgery. During the process, heart rate and mean arterial pressure were documented every 15 minutes. Numeric rating scale (NRS) for postoperative pain was recorded just after admission to the post-anesthesia care unit (PACU). If NRS was greater than 3, a 2.5-mg bolus dose of morphine sulfate was administered and repeated according to further pain assessments. Also, shivering assessment scale (BSAS) and post-

operative nausea and vomiting (PONV) score on a 4-point verbal descriptive scale were utilized to evaluate shivering and nausea status of patients in the PACU (11, 12).

2.2. Biochemical Method

Venous blood samples (5 mL each) were collected right before anesthesia induction (T0) and immediately after entering PACU after surgery and emergence (T1). After collecting the sample, the serum was allowed to clot for 5 - 10 minutes at room temperature. Then, the sample was centrifuged at 500 - 1000 G-force for 20 minutes, and the supernatants were collected carefully. Next, the antioxidants activity levels were analyzed using Zellbio (GmbH, Germany) kits. GPX, SOD, and CAT enzyme activities were measured in both T0 and T1 according to instructions provided by the kit manufacturer (data S1, S2, and S3).

2.3. Randomization and Blinding

Randomization was done using a random number table generated by computer. Participants were randomly assigned into Dex or control groups by simple randomization procedures (computerized random numbers). Since the intervention applied after induction of anesthesia, patients were blinded to the study. In addition, the staff that processed the samples and measured the desired items were blinded to study groups and samples.

2.4. Statistical Method

Sample size was calculated based on previous studies (13-15) by considering the comparison means and using sample size calculation formula with G power application.

In this study, all quantitative variables were presented as mean ± SD, and the categorical variables were shown as percentage. Also, some ordinal variables were shown as median with IQR (inter quartile range) in different groups.

To compare continuous variables between the study groups, the Student's *t*-test or non-parametric Mann-Whitney test was used alternatively. The categorical variables were compared using the Pearson chi-square test or Fisher's exact test when appropriate.

To compare the longitudinal data (heart rate and mean arterial pressure) between groups, the mixed model analysis was performed.

All the statistical tests were two-tailed at the significance level of 5%, and SPSS 26 was used for data analysis.

2.5. Ethics

The Ethical Committee of Shahid Beheshti University of medical sciences, Tehran, Iran approved this study, and a clinical trial code was obtained IRCT20151012024493N4. Also, written informed consent was obtained from all participants. The procedures were in accordance with the Helsinki Declaration of 1975, as revised in 2000.

3. Results

The C and D patients were not different in terms of gender, age, BMI, and concomitant diseases (Table 1).

Although the serum levels of CAT and GPX increased during the procedure, these changes were not statistically significant (P-value 0.579 and 0.762 respectively). SOD mean did not change over time in any of the groups. Also, SOD measures were not meaningfully different between the groups at any of the predetermined times (Table 2).

Three clinical outcomes were statistically different between the groups: (1) propofol consumption; (2) postoperative pain scores; and (3) shivering scores (Table 3). The duration of surgery and the volume of blood loss were not meaningfully different between the groups.

The amount of propofol consumed in DEX group was obviously less than control group, despite the similar surgery duration (1351.60 ± 427.17 mg in DEX group vs. 1715.80 ± 429.17 mg in control group, P-value = 0.007).

Pain scores measured in PACU were significantly less in DEX group (P-value < 0.001). The mean score reported for DEX group was 2.40 compared to 5.80 in the control group.

Based on Bedside Shivering Scale measures (0-3), the median score was 0 for DEX group and 1 for the control. This data was statistically interpreted in the form of interquartile range (IQR) and the difference was not significant (Table 3).

The heart rate and mean arterial pressure were measured in each time point for every patient and compared between the groups. The longitudinal changes over time were analyzed through mixed model analysis (Table 4). These measures were significantly lower in DEX group (P-value < 0.001). The visual demonstration of these data is presented in Figures 1 and 2.

4. Discussion

It has been demonstrated that different surgical procedures have strong impact on stress related responses (16). Other than surgery itself, several medical and non-medical processes could affect the inflammatory stress response, including various anesthesia medications and techniques (17-22).

On cellular level, exposure to volatile anesthetics manifested antioxidant effects on several cell types, except the neurons (23). Propofol has also proved to be a neuroprotective agent via its scavenging and immunomodulatory activities (24).

Desirable anti-inflammatory effects of DEX as a sedative in critical units are well known (25). In addition, perioperative adjunctive use of DEX in general anesthesia has been shown to considerably decrease serum IL6, IL8, and TNF- α level (26, 27). Furthermore, DEX blunts stress

responses via several other pathways, such as attenuating norepinephrine (28), epinephrine (29), and cortisol (27) release during surgical procedures. One of several known pathways for anesthesia and surgery-related oxidative stress is via attenuating antioxidants, such as GPX, SOD, and CAT (30). In a recent study, in patients with mild to moderate concomitant brain injuries who underwent neurologic surgeries, DEX ($1 \mu\text{g.kg}^{-1}.\text{hr}^{-1}$) was infused for 15 minutes before tracheal intubation, which demonstrated to have neuroprotective, anti-inflammatory, and antioxidant effects. Plasma brain derived neurotrophic factor (BDNF), several cytokines, and SOD levels were indicators of the above properties, respectively. SOD levels were significantly higher in patients receiving DEX compared to control group (31). In 2015, Han et al. compared three groups of DEX, propofol, and midazolam; each group received the anesthetic as induction and maintenance agent (13). Venous blood samples were obtained prior to the surgery (T0) and 2 h (T1), and 24 h (T2) after the surgery. At T1, all three markers decreased, and then decreased to less than the baseline values at T2.

On the other hand, the hemodynamic effects of DEX are documented repeatedly, which is also true in our study. Shivering and pain controlling effects are also compatible with other studies. In our prior study in transsphenoidal resection of pituitary adenoma, DEX infusion improved surgeon's satisfaction and decreased surgical site bleeding (32). However, current study did not indicate significant difference between groups regarding surgical blood loss.

SOD, GPX, and CAT play major roles as enzymatic antioxidant defense systems scavenging free oxygen radicals, mainly by reducing them into water. Inconsistent data about DEX effects on surgical stress responses have been reported. Various pathways have been assessed, and distinct surgical procedures may impact those responses or pathways differently. In the present study, particularly in laminectomy surgery, DEX did not affect the assessed antioxidants significantly. Further studies are recommended to clarify the possible mechanisms of observed discrepancies between various studies. Based on our findings, adjoining DEX to total intravenous anesthesia for simple spine surgeries may not affect antioxidant mechanisms significantly.

Equivocal findings in our study may be related to the limited number of cases. Therefore, larger studies with multiple measurements over an extended period of time could manifest more specific results. However, it should be noted that there are differences in the types of surgeries and their related stress responses. Laminectomy without dural puncture is a restricted source of oxidative stress compared to other neurosurgical procedures, and the results may vary substantially during other surgeries.

In conclusion, despite favorable hemodynamic and

Table 1. Demographic Characteristics of Participants

Variables	Control Group	Dex Group	P-Value
Male/ Female	15 (57.7)/11 (42.3)	15 (60.0)/10 (40.0)	0.867
Age (years)	43.88 ± 12.78	48.68 ± 11.79	0.144
BMI* (kg.m-2)	27.47 ± 3.47	26.77 ± 2.13	0.102
Concomitant Diseases	4 (15.4)	0 (0.0)	0.110
Total	26	25	NA

Abbreviation: BMI, body mass index.

Table 2. Serum Levels of Antioxidants Sampled Before Induction of Anesthesia and Immediately After Surgery in the Control and Dexmedetomidine Groups

Variables	Control Group	Dex Group	P-Value
CAT (U/mL)			
Before	2.57 ± 1.81	3.42 ± 3.61	0.935
After	4.89 ± 5.44	7.41 ± 14.14	0.607
Diff.	2.22 ± 6.33	3.98 ± 15.30	0.579
GPX (U/mL)			
Before	170.22 ± 116.81	131.66 ± 102.28	0.260
After	258.68 ± 165.51	238.23 ± 170.41	0.786
Diff.	88.45 ± 198.19	106.57 ± 176.95	0.762
SOD (U/mL)			
Before	82.49 ± 19.79	83.62 ± 15.24	0.860
After	84.24 ± 17.85	82.43 ± 27.60	0.567
Diff.	1.75 ± 30.26	-1.19 ± 35.01	0.665

Abbreviations: CAT, catalase enzyme activity; GPX, glutathione peroxidase enzyme activity; SOD, superoxide dismutase enzyme activity.

Table 3. Comparison of Clinical Outcomes Between the Control and Dexmedetomidine Groups

Variables	Control Group	Dex Group	P-Value
Duration of surgery (minutes)	165.20 ± 34.73	178.20 ± 51.35	0.318
Bleeding volume (mL)	365.25 ± 133.88	444.80 ± 282.36	0.810
Propofol consumption (mg)	1715.80 ± 429.17	1351.60 ± 427.17	0.007
NRS (0 - 10)	5.80 ± 1.00	2.40 ± 0.96	< 0.001
PONV Score; Median (IQR)	0 (0 - 0.75)	0 (0 - 0)	0.242
BSAS (0 - 3); Median (IQR)	1 (0 - 2)	0 (0 - 1)	0.030

Abbreviations: NRS, Numeric Rating Scale; PONV, postoperative nausea and vomiting; BSAS, Bedside Shivering Assessment Scale; IQR, inter quartile range.

Table 4. Mean Heart Rates (HR) and Mean Arterial Pressures (MAP) Changes over Time During the Surgery According to Mixed Model Analysis

Variables	Control Group	Dex Group	P-Value ^a
△ Heart rate	78.41 ± 12.16	68.32 ± 11.23	< 0.001
△ Mean arterial pressure	96.53 ± 15.70	83.61 ± 14.13	< 0.001

^a According to mixed model analysis

Figure 1. Heart rate during the procedure: comparison of mean heart rate measures during the surgery between the control and dexmedetomidine groups (error bars: ± 1 SE).

Figure 2. Mean arterial pressure during the procedure: comparison of mean arterial pressure measure during the surgery between the control and dexmedetomidine groups (error bars: ± 1 SE).

analgesic effects of dexmedetomidine, adjuvant infusion of this anesthetic during general anesthesia for laminectomy does not affect endogenous antioxidants during the surgery. Effects of the dexmedetomidine on antioxidant profile during postoperative period or in other spinal procedures are recommended for further studies.

Footnotes

Authors' Contribution: KM contributed to the concept, design, and definition of intellectual content, literature search, data analysis, statistical analysis, manuscript preparation, editing, and review. MN contributed to the

concept, design, and definition of intellectual content, data acquisition, and manuscript preparation. FS contributed to design, data acquisition, manuscript review and was the guarantor. SA and MB contributed to the concept, literature search, and data acquisition. PS contributed to concept, design, and definition of intellectual content, literature search, data acquisition, data analysis, statistical analysis, manuscript preparation, editing, and review.

Clinical Trial Registration Code:
IRCT20151012024493N4, 2019-03-13,
<https://www.irct.ir/trial/31328>

Conflict of Interests: The authors declare that there is no conflict of interest.

Data Reproducibility: The data presented in this study are openly available in one of the repositories or will be available on request from the corresponding author by this journal representative at any time during submission or after publication. Otherwise, all consequences of possible withdrawal or future retraction will be with the corresponding author.

Ethical Approval: The procedures were in accordance with the Helsinki Declaration of 1975, as revised in 2000 (IR.SBMU.RETECH.REC.1397.1029).

Funding/Support: The laboratory tests performed in this study were funded by Anesthesiology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Informed Consent: Written informed consent was obtained from all participants.

References

- Marana E, Annetta MG, Meo F, Pargaglioni R, Galeone M, Maussier ML, et al. Sevoflurane improves the neuroendocrine stress response during laparoscopic pelvic surgery. *Can J Anaesth*. 2003;50(4):348-54. doi: [10.1007/BF03021031](https://doi.org/10.1007/BF03021031). [PubMed: [12670811](https://pubmed.ncbi.nlm.nih.gov/12670811/)].
- Kuyumcu F, Aycan A. Evaluation of Oxidative Stress Levels and Antioxidant Enzyme Activities in Burst Fractures. *Med Sci Monit*. 2018;24:225-34. doi: [10.12659/msm.908312](https://doi.org/10.12659/msm.908312). [PubMed: [29324724](https://pubmed.ncbi.nlm.nih.gov/29324724/)]. [PubMed Central: [PMC5772452](https://pubmed.ncbi.nlm.nih.gov/PMC5772452/)].
- Cui J, Zhao H, Wang C, Sun JJ, Lu K, Ma D. Dexmedetomidine attenuates oxidative stress induced lung alveolar epithelial cell apoptosis in vitro. *Oxid Med Cell Longev*. 2015;2015:358396. doi: [10.1155/2015/358396](https://doi.org/10.1155/2015/358396). [PubMed: [25838866](https://pubmed.ncbi.nlm.nih.gov/25838866/)]. [PubMed Central: [PMC4369905](https://pubmed.ncbi.nlm.nih.gov/PMC4369905/)].
- Anupriya J, Kurhekar P. Randomised Comparison between the Efficacy of Two Doses of Nebulised Dexmedetomidine for Premedication in Paediatric Patients. *Turk J Anaesthesiol Reanim*. 2020;48(4):314-20. doi: [10.5152/TJAR.2019.78889](https://doi.org/10.5152/TJAR.2019.78889). [PubMed: [32864647](https://pubmed.ncbi.nlm.nih.gov/32864647/)]. [PubMed Central: [PMC7434347](https://pubmed.ncbi.nlm.nih.gov/PMC7434347/)].
- Srivastava VK, Agrawal S, Kumar S, Khan S, Sharma S, Kumar R. Comparative Evaluation of Dexmedetomidine and Propofol Along With Scalp Block on Haemodynamic and Postoperative Recovery for Chronic Subdural Haematoma Evacuation Under Monitored Anaesthesia Care. *Turk J Anaesthesiol Reanim*. 2018;46(1):51-6. doi: [10.5152/TJAR.2018.16878](https://doi.org/10.5152/TJAR.2018.16878). [PubMed: [30140501](https://pubmed.ncbi.nlm.nih.gov/30140501/)]. [PubMed Central: [PMC5858890](https://pubmed.ncbi.nlm.nih.gov/PMC5858890/)].
- Suvvari P, Mishra S, Bhatnagar S, Garg R, Bharati SJ, Gupta N, et al. Comparison of Intranasal Dexmedetomidine Versus Intranasal Ketamine as Premedication for Level of Sedation in Children Undergoing Radiation Therapy: A Prospective, Randomised, Double-Blind Study. *Turk J Anaesthesiol Reanim*. 2020;48(3):215-22. doi: [10.5152/TJAR.2019.45087](https://doi.org/10.5152/TJAR.2019.45087). [PubMed: [32551449](https://pubmed.ncbi.nlm.nih.gov/32551449/)]. [PubMed Central: [PMC7279870](https://pubmed.ncbi.nlm.nih.gov/PMC7279870/)].
- Arpaci AH, Bozkirli F. Comparison of sedation effectiveness of remifentanyl-dexmedetomidine and remifentanyl-midazolam combinations and their effects on postoperative cognitive functions in cystoscopies: A randomized clinical trial. *J Res Med Sci*. 2013;18(2):107-14. [PubMed: [23914211](https://pubmed.ncbi.nlm.nih.gov/23914211/)]. [PubMed Central: [PMC3724369](https://pubmed.ncbi.nlm.nih.gov/PMC3724369/)].
- Turkkan M, Gulec E, Hatipoglu Z, Ilginel MT, Ozcengiz D. The Effect of Sevoflurane and Dexmedetomidine on Pulmonary Mechanics in ICU Patients. *Turk J Anaesthesiol Reanim*. 2019;47(3):206-12. doi: [10.5152/TJAR.2019.37108](https://doi.org/10.5152/TJAR.2019.37108). [PubMed: [31183467](https://pubmed.ncbi.nlm.nih.gov/31183467/)]. [PubMed Central: [PMC6537958](https://pubmed.ncbi.nlm.nih.gov/PMC6537958/)].
- Uysal HY, Tezer E, Turkoglu M, Aslanargun P, Basar H. The effects of dexmedetomidine on hemodynamic responses to tracheal intubation in hypertensive patients: A comparison with esmolol and sufentanil. *J Res Med Sci*. 2012;17(1):22-31. [PubMed: [23248653](https://pubmed.ncbi.nlm.nih.gov/23248653/)]. [PubMed Central: [PMC3523434](https://pubmed.ncbi.nlm.nih.gov/PMC3523434/)].
- Tang P, Zhu R, Gu Y, Gu JM, Xie ZA, Xu JQ, et al. Ascorbic Acid Attenuates Multifidus Muscles Injury and Atrophy After Posterior Lumbar Spine Surgery by Suppressing Inflammation and Oxidative Stress in a Rat Model. *Spine (Phila Pa 1976)*. 2018;43(21):E1249-59. doi: [10.1097/BRS.0000000000002669](https://doi.org/10.1097/BRS.0000000000002669). [PubMed: [29649092](https://pubmed.ncbi.nlm.nih.gov/29649092/)].
- Badjatia N. Shivering: scores and protocols. *Crit Care*. 2012;16(S2). doi: [10.1186/cc11267](https://doi.org/10.1186/cc11267).
- Boogaerts JG, Vanacker E, Seidel L, Albert A, Bardiau FM. Assessment of postoperative nausea using a visual analogue scale. *Acta Anaesthesiol Scand*. 2000;44(4):470-4. doi: [10.1034/j.1399-6576.2000.440420.x](https://doi.org/10.1034/j.1399-6576.2000.440420.x). [PubMed: [10757584](https://pubmed.ncbi.nlm.nih.gov/10757584/)].
- Han C, Ding W, Jiang W, Chen YU, Hang D, Gu D, et al. A comparison of the effects of midazolam, propofol and dexmedetomidine on the antioxidant system: A randomized trial. *Exp Ther Med*. 2015;9(6):2293-8. doi: [10.3892/etm.2015.2410](https://doi.org/10.3892/etm.2015.2410). [PubMed: [26136976](https://pubmed.ncbi.nlm.nih.gov/26136976/)]. [PubMed Central: [PMC4473377](https://pubmed.ncbi.nlm.nih.gov/PMC4473377/)].
- Khoshraftar E, Ranjbar A, Kharkhane B, Tavakol Heidary S, Gharebaghi Z, Zadkhosh N. Antioxidative effects of propofol vs. ketamin in individuals undergoing surgery. *Arch Iran Med*. 2014;17(7):486-9. [PubMed: [24979561](https://pubmed.ncbi.nlm.nih.gov/24979561/)].
- Bostankolu E, Ayoglu H, Yurtlu S, Okyay RD, Erdogan G, Deniz Y, et al. Dexmedetomidine did not reduce the effects of tourniquet-induced ischemia-reperfusion injury during general anesthesia. *Kaohsiung J Med Sci*. 2013;29(2):75-81. doi: [10.1016/j.kjms.2012.08.013](https://doi.org/10.1016/j.kjms.2012.08.013). [PubMed: [23347808](https://pubmed.ncbi.nlm.nih.gov/23347808/)].
- Crippa J, Mari GM, Miranda A, Costanzi AT, Maggioni D. Surgical Stress Response and Enhanced Recovery after Laparoscopic Surgery - A systematic review. *Chirurgia (Bucur)*. 2018;113(4):455-63. doi: [10.21614/chirurgia.113.4.455](https://doi.org/10.21614/chirurgia.113.4.455). [PubMed: [30183575](https://pubmed.ncbi.nlm.nih.gov/30183575/)].
- Fu VX, Oomens P, Smeiders D, van den Berg SAA, Feelders RA, Wijnhoven BPL, et al. The Effect of Perioperative Music on the Stress Response to Surgery: A Meta-analysis. *J Surg Res*. 2019;244:444-55. doi: [10.1016/j.jss.2019.06.052](https://doi.org/10.1016/j.jss.2019.06.052). [PubMed: [31326711](https://pubmed.ncbi.nlm.nih.gov/31326711/)].
- Brodner G, Van Aken H, Hertle L, Fobker M, Von Eckardstein A, Goeters C, et al. Multimodal perioperative management-combining thoracic epidural analgesia, forced mobilization, and oral nutrition-reduces hormonal and metabolic stress and improves convalescence after major urologic surgery. *Anesth Analg*. 2001;92(6):1594-600. doi: [10.1097/0000539-200106000-00049](https://doi.org/10.1097/0000539-200106000-00049). [PubMed: [11375853](https://pubmed.ncbi.nlm.nih.gov/11375853/)].
- Vosoughian M, Dahi-Taleghani M, Moshari M, Rajaei S, Rajabi S, Taheri F. The Effect of Spinal and General Anesthesia on Cytokine Serum Levels Following Cesarean Section in Preeclampsia. *J Cell Mol Anesth*. 2018;3(3):89-97.
- Faani K, Shadnough M, Jahangirifard A, Foroughi M, Jelveh-Moghaddam HA, Dabbagh A. The effect of 1, 25 (OH) 2 Vitamin D3 on serum levels of inflammatory cytokines in patients undergoing CABG. *J Cell Mol Anesth*. 2018;3(3):98-102.

21. Pi J, Sun Y, Zhang Z, Wan C. Combined anesthesia shows better curative effect and less perioperative neuroendocrine disorder than general anesthesia in early stage NSCLC patients. *J Int Med Res.* 2019;**47**(10):4743–52. doi: [10.1177/0300060519862102](https://doi.org/10.1177/0300060519862102). [PubMed: [31510831](https://pubmed.ncbi.nlm.nih.gov/31510831/)]. [PubMed Central: [PMC6833388](https://pubmed.ncbi.nlm.nih.gov/PMC6833388/)].
22. Karabayirli S, Keskin EA, Kaya A, Koca C, Erel O, Demircioglu RI, et al. Assessment of fetal antioxidant and oxidant status during different anesthesia techniques for elective cesarean sections. *J Res Med Sci.* 2015;**20**(8):739–44. doi: [10.4103/1735-1995.168373](https://doi.org/10.4103/1735-1995.168373). [PubMed: [26664420](https://pubmed.ncbi.nlm.nih.gov/26664420/)]. [PubMed Central: [PMC4652306](https://pubmed.ncbi.nlm.nih.gov/PMC4652306/)].
23. Lee YM, Song BC, Yeum KJ. Impact of Volatile Anesthetics on Oxidative Stress and Inflammation. *Biomed Res Int.* 2015;**2015**:242709. doi: [10.1155/2015/242709](https://doi.org/10.1155/2015/242709). [PubMed: [26101769](https://pubmed.ncbi.nlm.nih.gov/26101769/)]. [PubMed Central: [PMC4458520](https://pubmed.ncbi.nlm.nih.gov/PMC4458520/)].
24. Kotani Y, Shimazawa M, Yoshimura S, Iwama T, Hara H. The experimental and clinical pharmacology of propofol, an anesthetic agent with neuroprotective properties. *CNS Neurosci Ther.* 2008;**14**(2):95–106. doi: [10.1111/j.1527-3458.2008.00043.x](https://doi.org/10.1111/j.1527-3458.2008.00043.x). [PubMed: [18482023](https://pubmed.ncbi.nlm.nih.gov/18482023/)]. [PubMed Central: [PMC6494023](https://pubmed.ncbi.nlm.nih.gov/PMC6494023/)].
25. Ohta Y, Miyamoto K, Kawazoe Y, Yamamura H, Morimoto T. Effect of dexmedetomidine on inflammation in patients with sepsis requiring mechanical ventilation: a sub-analysis of a multicenter randomized clinical trial. *Crit Care.* 2020;**24**(1):493. doi: [10.1186/s13054-020-03207-8](https://doi.org/10.1186/s13054-020-03207-8). [PubMed: [32778146](https://pubmed.ncbi.nlm.nih.gov/32778146/)]. [PubMed Central: [PMC7416813](https://pubmed.ncbi.nlm.nih.gov/PMC7416813/)].
26. Li B, Li Y, Tian S, Wang H, Wu H, Zhang A, et al. Anti-inflammatory Effects of Perioperative Dexmedetomidine Administered as an Adjunct to General Anesthesia: A Meta-analysis. *Sci Rep.* 2015;**5**:12342. doi: [10.1038/srep12342](https://doi.org/10.1038/srep12342). [PubMed: [26196332](https://pubmed.ncbi.nlm.nih.gov/26196332/)]. [PubMed Central: [PMC4508837](https://pubmed.ncbi.nlm.nih.gov/PMC4508837/)].
27. Jain A, Sinha R, Pandey S, Sahu V. Comparative Evaluation of Dexmedetomidine and Pregabalin as Premedication Agent to Attenuate Adverse Hemodynamic and Stress Response in Patients Undergoing Laparoscopic Cholecystectomy. *Anesth Essays Res.* 2019;**13**(4):608–14. doi: [10.4103/aer.AER_115_19](https://doi.org/10.4103/aer.AER_115_19). [PubMed: [32009703](https://pubmed.ncbi.nlm.nih.gov/32009703/)]. [PubMed Central: [PMC6937902](https://pubmed.ncbi.nlm.nih.gov/PMC6937902/)].
28. Kang R, Jeong JS, Ko JS, Lee SY, Lee JH, Choi SJ, et al. Intraoperative dexmedetomidine attenuates norepinephrine levels in patients undergoing transsphenoidal surgery: a randomized, placebo-controlled trial. *BMC Anesthesiol.* 2020;**20**(1):100. doi: [10.1186/s12871-020-01025-7](https://doi.org/10.1186/s12871-020-01025-7). [PubMed: [32359367](https://pubmed.ncbi.nlm.nih.gov/32359367/)]. [PubMed Central: [PMC7195722](https://pubmed.ncbi.nlm.nih.gov/PMC7195722/)].
29. Kim MH, Lee KY, Bae SJ, Jo M, Cho JS. Intraoperative dexmedetomidine attenuates stress responses in patients undergoing major spine surgery. *Minerva Anesthesiol.* 2019;**85**(5):468–77. doi: [10.23736/S0375-9393.18.12992-0](https://doi.org/10.23736/S0375-9393.18.12992-0). [PubMed: [30226342](https://pubmed.ncbi.nlm.nih.gov/30226342/)].
30. Zhang J, Gao J, Guo G, Li S, Zhan G, Xie Z, et al. Anesthesia and surgery induce delirium-like behavior in susceptible mice: the role of oxidative stress. *Am J Transl Res.* 2018;**10**(8):2435–44. [PubMed: [30210682](https://pubmed.ncbi.nlm.nih.gov/30210682/)]. [PubMed Central: [PMC6129548](https://pubmed.ncbi.nlm.nih.gov/PMC6129548/)].
31. Ding Q, Zhang X, Chen P. Intraoperative Dexmedetomidine in Peripheral or Emergency Neurologic Surgeries of Patients With Mild-to-Moderate Traumatic Brain Injuries: A Retrospective Cohort Study. *Dose Response.* 2020;**18**(2):1559325820920120. doi: [10.1177/1559325820920119](https://doi.org/10.1177/1559325820920119). [PubMed: [32489336](https://pubmed.ncbi.nlm.nih.gov/32489336/)]. [PubMed Central: [PMC7238827](https://pubmed.ncbi.nlm.nih.gov/PMC7238827/)].
32. Salimi A, Sharifi G, Bahrani H, Mohajerani SA, Jafari A, Safari F, et al. Dexmedetomidine could enhance surgical satisfaction in Trans-sphenoidal resection of pituitary adenoma. *J Neurosurg Sci.* 2017;**61**(1):46–52. doi: [10.23736/S0390-5616.16.02792-2](https://doi.org/10.23736/S0390-5616.16.02792-2). [PubMed: [24866894](https://pubmed.ncbi.nlm.nih.gov/24866894/)].