



The Influential Factors in Burnout in the Healthcare Staff in Kermanshah, Iran

Mostafa Jafari^{1*}, Sorya Karimi², Leila Rahmani², Farshad Nadri² and Mansour Rezaei²

¹Health Faculty, Kermanshah University of Medical Sciences, Kermanshah, Iran

²Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding author: Health Faculty, Kermanshah University of Medical Sciences, Kermanshah, Iran. Email: mostafajafari66@yahoo.com

Received 2021 June 29; Revised 2021 August 30; Accepted 2021 October 12.

Abstract

Background: Burnout plays a key role in the life and physical and mental health of individuals and may reduce the efficiency and motivation of healthcare staff.

Objectives: The present study aimed to evaluate the influential factors in burnout in the healthcare staff of Kermanshah, Iran.

Methods: This cross-sectional study was conducted on the healthcare staff in Kermanshah in 2019. Data were collected using a demographic questionnaire and the Maslach burnout inventory (Cronbach's alpha: 0.78). The participants were selected via census sampling (n = 282). Data analysis was performed in SPSS version 22 using the Kolmogorov-Smirnov test, Mann-Whitney U test, Kruskal-Wallis test, and Spearman's correlation-coefficient.

Results: Significant correlations were observed between emotional fatigue, disease history (P = 0.006), and financial problems (P = 0.05), as well as between depersonalization and financial difficulties (P = 0.024) and between personal adequacy and family disease history (P = 0.015). However, no significant associations between burnout and gender, educational level, professional status, marital status and living situation were shown.

Conclusions: Considering the high prevalence of burnout in the healthcare staff with a disease history in themselves and their families and the association between burnout and financial problems, special attention should be paid to the health status of healthcare status through measures such as stress management and counseling to diminish occupational stress and the impact of burnout on these employees. In addition, measures such as the implementation of leisure and educational programs and stress management workshops are recommended.

Keywords: Burnout, Depersonalization, Personal Adequacy, Emotional Fatigue

1. Background

Burnout is defined as mental and physical fatigue caused by high and chronic stress. If the stress continues, the motivation and vitality of the individual will decrease significantly. Burnout reduces energy and feelings of hope, strength, and capability, while reinforcing feelings of pessimism and resentment. Job dissatisfaction due to burnout could lead to job loss, the disappearance of social relations, and even health issues (1).

The phenomenon of burnout caused by workplace pressures and type of work has been observed in 40% of the population in industrial societies (2). Medical professionals are also faced with this issue depending on the type of their profession. Stress caused by patient care may have long-term debilitating effects on healthcare staff (3). According to the studies conducted in Iran, burnout is highly common among nurses. A study of the nurses in a hospi-

tal in Kermanshah (Iran) indicated that 39.2% of the nurses had high levels of emotional fatigue, 37.6% had significant depersonalization, and 73.5% had low personal adequacy (4). In another study conducted on the healthcare staff in Babol (Iran), Hosseini et al. reported that 46 employees (22.2%) had significant emotional exhaustion, and 55 subjects (26.6%) had high depersonalization. However, none of the respondents experienced low personal adequacy (5).

According to the literature, burnout could interfere with proper task management in the workplace, cause stress, and change the health status of nurses (6); therefore, special attention should be paid to this issue in order to eliminate the root causes. The issue of burnout in healthcare staff has been investigated in different countries, including Iran. In a study conducted in Taiwan, the prevalence of burnout in healthcare providers was assessed based on factors such as youth, full-time job, being a nurse or physician assistant, employment status, and so-

cial commitment (7).

In addition to nurses, midwives seem to be at a high risk of burnout due to the type of their profession. According to a study conducted in Urmia (Iran), a quarter of midwives suffered from extreme emotional fatigue, and more than half of these subjects experienced moderate-to-high levels of emotional fatigue (8). Research also suggests that constant exposure to patients, being held accountable for human health, performing clinical procedures, and managing frequent emergencies are among the main factors that reduce the quality of health services and contribute to the depression and fatigue of healthcare providers (9).

Previous findings have indicated a strong correlation between burnout and reduced productivity in organizations (10). The most important issues caused by burnout are absenteeism, leaving the service, consecutive delays, job changes, and reduced quality of patient care (11). In addition, the effect of burnout on job satisfaction has been confirmed, indicating a strong correlation between emotional fatigue and job satisfaction. Therefore, organizations must attempt to minimize burnout in their employees through preventive measures such as stress management and counseling with employees to manage occupational stress and lower the impact of burnout on the employees (12).

2. Objectives

Since no studies have investigated burnout in the healthcare staff in Kermanshah, the present study aimed to evaluate the influential factors in the burnout of these healthcare professionals.

3. Methods

This cross-sectional descriptive-analytical study, carried out in Kermanshah, Iran, in 2019, included 282 Kermanshah health center employees who were selected through census samples. Participants included doctors, midwives, nurses, health professionals and administrative staff.

The data were collected using a demographic questionnaire consisting of 12 items on age, gender, level of education, work experience, marital status, number of children, employment status, medical history, family history, economic situation, economic problems and the situation of the place of residence.

3.1. Maslach Burnout Inventory

Another instrument used in this study was the Maslach burnout inventory, which consists of 22 statements to measure three aspects of burnout, nine statements to assess

emotional fatigue, five statements to evaluate depersonalization, and eight statements regarding personal adequacy. The propositions were scored based on a seven-point Likert scale (never = 0, very low = 1, low = 2, moderate = 3, high = 4, very high = 5, extremely high = 6) (13). The main dimensions of the questionnaire were emotional fatigue, depersonalization, and personal adequacy.

For the first time in Iran, the validity and reliability of this inventory were confirmed by Moalemi, and the internal compatibility coefficient was also confirmed at the Cronbach's alpha of 0.78 (14). Furthermore, Sadr Khan Lou et al. estimated the reliability coefficient of this inventory (Cronbach's alpha: 0.78), and the Cronbach's alpha coefficient of the entire scale and its three dimensions has been reported to be 0.65 - 0.85 (8).

In the present study, the researcher initially explained the objectives of the study and instructed the participants on completing the instruments. In addition, the participants were assured of confidentiality terms regarding their personal information. After obtaining informed consent, the questionnaires were provided to the participants to be completed.

Data analysis was performed in SPSS version 22 using the Kolmogorov-Smirnov (KS) test to determine the normality of the data. The KS test results indicated that the distribution of the variables was nonparametric and the Mann-Whitney U test, Kruskal-Wallis test, and Spearman's correlation-coefficient were applied to assess the differences or correlations between the variables. In all the statistical analyses, the significance level was set at 0.05.

4. Results

In total, 282 healthcare providers in Kermanshah were surveyed, including 72 and 28% female and male staff, respectively. The results of various variables and the statistical relationships between them are summarized in Tables 1 and 2. According to tables results significant correlations were observed between emotional fatigue and disease history ($P = 0.006$), financial problems ($P = 0.05$), also personal adequacy and family disease history ($P = 0.015$), and then between depersonalization and financial problems ($P = 0.024$).

5. Discussion

The results of the present study indicated no significant correlations between burnout, age, and marital status. Emotional fatigue and personal adequacy were observed to be more significant in the married subjects compared to the single and divorced subjects, while depersonalization was more significant in the divorced subjects

Table 1. Comparison of the Mean (\pm Standard Deviation) of Burnout Dimension Variables with the Studied Variables

Variables	Dimensions of Burnout			Burnout
	Emotional Fatigue	Depersonalization	Personal Adequacy	
Gender				
Female	42.76 ± 9.749	27.66 ± 2.871	29.81 ± 8.175	100.24 ± 16.435
Male	43.08 ± 10.103	26.70 ± 3.639	27.75 ± 9.557	97.52 ± 16.914
P-value	0.723	0.060	0.071	0.218
Level of Education				
Diploma and less	47.087 ± 7.097	25.547 ± 3.676	27.74 ± 8.864	99.76 ± 18.282
Associate Degree	42.84 ± 9.292	28.12 ± 2.447	29.39 ± 9.374	100.35 ± 14.839
Bachelor	42.45 ± 9.993	27.33 ± 3.114	29.18 ± 8.174	99.08 ± 16.557
Master's degree and higher	43.17 ± 10.096	26.72 ± 3.453	30.47 ± 8.564	100.79 ± 18.163
P-value	0.788	0.413	0.342	0.775
Staff Status				
Official	38.895 ± 4.832	28.25 ± 1.512	29.39 ± 8.72	99.66 ± 16.182
Contractual	43.74 ± 9.937	27.39 ± 39.448	28.09 ± 8.974	99.22 ± 17.904
Corporate and service contract	43.56 ± 11.260	26.42 ± 3.639	28.87 ± 7.785	98.20 ± 18.985
P-value	0.797	0.448	0.792	0.924
Marital status				
Single	42.53 ± 10.968	27.643 ± 3.463	29.07 ± 8.698	99.71 ± 16.905
Married	43.00 ± 9.555	27.37 ± 3.030	29.34 ± 8.855	99.63 ± 16.414
Divorced	41.585 ± 16.374	28.50 ± 2.598	27.00 ± 7.000	99.40 ± 24.704
P-value	0.291	0.215	0.767	0.866
History of disease				
Yes	39.27 ± 10.906	27.81 ± 2.679	29.00 ± 8.778	96.08 ± 16.948
No	43.66 ± 9.477	27.29 ± 3.223	29.28 ± 8.601	100.25 ± 16.442
P-value	0.006	0.503	0.831	0.102
History of disease in the family				
Yes	42.32 ± 9.477	27.67 ± 2.938	31.37 ± 7.239	101.16 ± 15.117
No	43.02 ± 9.964	27.30 ± 3.194	28.51 ± 8.939	98.90 ± 17.048
P-value	0.444	0.463	0.015	0.326
Financial problems				
Yes	40.90 ± 9.809	26.68 ± 3.402	29.52 ± 8.528	96.82 ± 15.846
No	43.35 ± 9.797	27.57 ± 3.035	29.21 ± 8.635	100.21 ± 16.783
P-value	0.05	0.024	0.750	0.113
Housing				
Rent	43.70 ± 10.517	26.90 ± 3.655	27.77 ± 9.208	98.38 ± 18.299
Proprietary	42.75 ± 9.416	27.61 ± 2.908	29.83 ± 8.536	100.20 ± 15.917
Other	40.88 ± 12.288	26.53 ± 3.502	27.35 ± 6.773	94.76 ± 17.967
P-value	0.505	0.253	0.187	0.558

Table 2. Comparison of Pearson Correlation Coefficient (r) Between Burnout Dimensions with Studied Variables

Variables	Emotional Fatigue		Depersonalization		Personal Adequacy		Burnout	
	r	P-value	r	P-value	r	P-value	r	P-value
Age	-0.005	0.934	0.031	0.604	-0.011	0.860	-0.012	0.845
Work experience	-0.039	0.516	0.078	0.194	-0.013	0.834	-0.021	0.727
Number of children	-0.076	0.263	-0.104	0.125	-0.020	0.764	-0.060	0.374
Income	-0.027	0.651	-0.020	0.733	-0.024	0.691	-0.024	0.693

compared to the single individuals. Our findings regarding the single and married subjects are consistent with the study by Sohrabi et al. (15).

According to the current research, the mean scores of the burnout dimensions had no significant differences between the male and female subjects. On the other hand, emotional fatigue was more significant in men compared to women, while depersonalization and personal adequacy were more significant in women compared to men. These differences could be attributed to factors such as emotional instability, high levels of stress, accuracy in performing professional duties, and more responsibility affecting the relations of the healthcare staff with patients. In the present study, emotional fatigue was observed in 29% of the subjects, depersonalization was denoted in 19%, and low personal adequacy was observed in 44%. In more than 33% of the participants, at least two of the burnout dimensions were considered significant (16).

According to our findings, work experience had no significant correlation with the mean scores of the dimensions of burnout. Emotional fatigue and personal adequacy were inversely correlated, while depersonalization was directly correlated with work experience. Increased work experience also led to the reduction of career excitement and personal adequacy over time, while depersonalization increased with more work experience; this is in line with the study by Mahmoudi et al. (17). Furthermore, no significant correlation was observed between burnout and academic degree in the present study, which is consistent with the findings of Sadr Khan Lou et al. regarding burnout in the midwives of the health centers in Urmia (8). The current study did not show any significant correlations between the dimensions burnout and employment status and income status, which corresponds to the study by Cavus and Demir (18).

In the present study, 18.4% of the participants had a disease history, and a significant correlation was observed between emotional fatigue and disease history. This could be attributed to factors such as stress, anxiety, and mood problems in a diseased state. Moreover, 25.2% of the participants had a family disease history, and a significant associ-

ation was denoted between the personal adequacy dimension and family disease history. Significant correlations were also denoted between financial problems and the dimensions of emotional fatigue and personal adequacy. In terms of the housing status, 72.3% of the participants were homeowners, 21.6% were tenants, and 6% had other statuses who did not provide any burnout statements.

5.1. Conclusion

According to the results, the dimensions of burnout were more significant in the employees with a disease history, family medical history, and financial problems. Therefore, it is suggested that measures such as implementing recreational and educational programs and holding stress management workshops be taken by health policymakers to increase the productivity and satisfaction of healthcare staff by providing a calm work environment.

Acknowledgments

The staff of Kermanshah health centers who have participated in this study is highly thanked and appreciated.

Footnotes

Authors' Contribution: Study concept and design, M J; Data analysis and interpretation, M R; Drafting of the manuscript, M J, S K and L R; Critical review of the manuscript for important intellectual content, M J and F N. All authors submitted comments and approved the final manuscript.

Conflict of Interests: There is no conflict of interest between the authors.

Funding/Support: This study was not funded by the KUMS Research deputy.

References

1. Nazari H, Jariani M, Beiranvand S, Saki M, Aghajeri N, Ebrahimzadeh F. The Prevalence of Job Stress and its Relationship with Burnout Syndrome among the Academic Members of Lorestan University of Medical Sciences. *J Caring Sci.* 2016;5(1):75-84. doi: [10.15171/jcs.2016.008](https://doi.org/10.15171/jcs.2016.008). [PubMed: [26989668](https://pubmed.ncbi.nlm.nih.gov/26989668/)]. [PubMed Central: [PMC4794547](https://pubmed.ncbi.nlm.nih.gov/PMC4794547/)].

2. Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol.* 2001;**52**:397–422. doi: [10.1146/annurev.psych.52.1.397](https://doi.org/10.1146/annurev.psych.52.1.397). [PubMed: [11148311](https://pubmed.ncbi.nlm.nih.gov/11148311/)].
3. Ebrahimi H, Navidian A, Ameri M, Sadeghi M. [Burnout, dimensions and its related factors in the operational staff of medicine emergency]. *J Promot Manag.* 2014;**3**(3):16–26. Persian.
4. Ziaei M, Yarmohammadi H, Karamimatin B, Yarmohammadi S, Nazari Z, Gharagozlou F. [Prevalence and risk factors of occupational burnout among nurses of a hospital in Kermanshah in 2013]. *Iran J Ergon.* 2014;**2**(2):67–74. Persian.
5. Hosseini S, Aghalari Z, Ashrafian Amiri H, Bayani F, Gholinia H, Jafarian S. [Occupational Burnout and its Related Factors among Health Care Center Staff in Babol]. *Health Dev J.* 2020;**8**(3):246–57. Persian.
6. Khamisa N, Peltzer K, Oldenburg B. Burnout in relation to specific contributing factors and health outcomes among nurses: A systematic review. *Int J Environ Res Public Health.* 2013;**10**(6):2214–40. doi: [10.3390/ijerph10062214](https://doi.org/10.3390/ijerph10062214). [PubMed: [23727902](https://pubmed.ncbi.nlm.nih.gov/23727902/)]. [PubMed Central: [PMC3717733](https://pubmed.ncbi.nlm.nih.gov/PMC3717733/)].
7. Chou LP, Li CY, Hu SC. Job stress and burnout in hospital employees: Comparisons of different medical professions in a regional hospital in Taiwan. *BMJ Open.* 2014;**4**(2). e004185. doi: [10.1136/bmjopen-2013-004185](https://doi.org/10.1136/bmjopen-2013-004185). [PubMed: [24568961](https://pubmed.ncbi.nlm.nih.gov/24568961/)]. [PubMed Central: [PMC3939670](https://pubmed.ncbi.nlm.nih.gov/PMC3939670/)].
8. Sadrkhanlou M, Ranjii A. [Occupational burnout among midwives working in health centers in Urmia and its relationship with their working status in 2010]. *Journal of North Khorasan University of Medical Sciences.* 2013;**5**(1):115–24. Persian. doi: [10.29252/jnkums.5.1.115](https://doi.org/10.29252/jnkums.5.1.115).
9. Zanganeh S, Moradbeygi K, Rasteh M, Raisifar Z. [Investigating the Degree of Job Burnout and Its Relation with the General Health of Nurses Working in Hospitals of Abadan and Khoramshahr in 2012]. *J Ilam Uni Med Sci.* 2014;**22**(Supl):217–24. Persian.
10. Dewa CS, Loong D, Bonato S, Thanh NX, Jacobs P. How does burnout affect physician productivity? A systematic literature review. *BMC Health Serv Res.* 2014;**14**:325. doi: [10.1186/1472-6963-14-325](https://doi.org/10.1186/1472-6963-14-325). [PubMed: [25066375](https://pubmed.ncbi.nlm.nih.gov/25066375/)]. [PubMed Central: [PMC4119057](https://pubmed.ncbi.nlm.nih.gov/PMC4119057/)].
11. Taei M, Divsalar K. Frequency of burnout among general practitioners of Kerman city, 2008. *J Kerman Univ Medical Sc.* 2010;**17**(3):268–76.
12. Eslami H, Amini A, Babaeian R, Poyeh M, Dehghan SM, Nasiri M. Assessment of Job Burnout Among Staff in Medical Diagnostic Laboratories in Shahid Sadoughi University of Medical Sciences in Yazd in 2013. *J Rafsanjan Univ Med Sci.* 2014;**13**(6):497–508.
13. Maslach C, Jackson SE. The measurement of experienced burnout. *J Organ Behav.* 1981;**2**(2):99–113. doi: [10.1002/job.4030020205](https://doi.org/10.1002/job.4030020205).
14. Moalemi S, Kavosi Z, Beygi N, Deghan A, Karimi A, Parvizi MM. Evaluation of the Persian Version of Maslach Burnout Inventory-Human Services Survey among Iranian Nurses: Validity and Reliability. *Galen Med J.* 2018;**7**. e995. doi: [10.22086/gmj.v0i0.995](https://doi.org/10.22086/gmj.v0i0.995). [PubMed: [34466422](https://pubmed.ncbi.nlm.nih.gov/34466422/)]. [PubMed Central: [PMC8343696](https://pubmed.ncbi.nlm.nih.gov/PMC8343696/)].
15. Sohrabi H, Majzobi S, Shahoei R. [Burnout rate and related factors among midwives]. *Sci J Nursing, Midwifery and Paramedical Faculty.* 2014;**3**(6):67–76. Persian.
16. De la Fuente-Solana EI, Suleiman-Martos N, Pradas-Hernandez L, Gomez-Urquiza JL, Canadas-De la Fuente GA, Albendin-Garcia L. Prevalence, Related Factors, and Levels of Burnout Syndrome Among Nurses Working in Gynecology and Obstetrics Services: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health.* 2019;**16**(14). doi: [10.3390/ijerph16142585](https://doi.org/10.3390/ijerph16142585). [PubMed: [31331046](https://pubmed.ncbi.nlm.nih.gov/31331046/)]. [PubMed Central: [PMC6678444](https://pubmed.ncbi.nlm.nih.gov/PMC6678444/)].
17. Mahmoudi G, Alhani F, QanbarRohi A. [Related about some factors to levels of burnout in nurses working in Golestan State]. *J Nurs Midwifery Care.* Persian.
18. Çavus MF, Demir Y. The Impacts of Structural and Psychological Empowerment on Burnout: A Research on Staff Nurses in Turkish State Hospitals Canadian Social science. *Can Soc Sci.* 2010;**6**(2):63–72. doi: [10.3968/j.css.1923669720100604.007](https://doi.org/10.3968/j.css.1923669720100604.007).