Relaxation Effects of Aromatherapy Intervention on Female and Male Anxiety: A Randomized Control Trial

Jamshid Eslami ¹, Abed Ebrahimi ², Isan Darvishi ³, Khadijeh Momeni ² and Marzieh Akbarzadeh ^{4, *}

¹Community Based Psychiatric Care Research Center, Department of Anesthesia and Surgical Technologists, School of Nursing and Midwifery, Shiraz University Medical Science, Shiraz, Iran

²Department of Operating Room, Faculty of Para Medicine, Bushehr University of Medical Sciences, Bushehr, Iran

³Department of Operating Room, School of Nursing and Midwifery, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁴Maternal-Fetal Medicine Research Center, Department of Midwifery, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding author: Maternal-Fetal Medicine Research Center, Department of Midwifery, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran. Email: akbarzadm@sums.ac.ir

Received 2020 November 23; Revised 2021 March 14; Accepted 2021 May 23.

Abstract

Background: The control or reduction of anxiety before surgery is one of the most important components of preoperative care. The question is whether anxiety differs in male and female candidates for surgery.

Objectives: The current study aimed to evaluate the effect of aromatherapy with *Citrus aurantium* on anxiety in men and women candidates for surgery.

Methods: This single-blind randomized clinical trial was performed on 60 patients and controlled by placebo. The patients were divided into 30 subjects in the control group and 30 subjects in the aromatherapy group (inhaling *C. aurantium* essential oil). The patients completed the Spielberger state-trait anxiety inventory before and after the intervention. After collecting information, the questionnaire information was analyzed using SPSS software (version 16).

Results: After aromatherapy, 20% of men had mild state and trait anxiety. Moreover, 80% of men had relatively mild state and trait anxiety. In the group of women, 20% of the cases had mild state and trait anxiety. Furthermore, 73.3% of women had relatively mild state and trait anxiety. Additionally, 6.7% of women had relatively severe anxiety. Anxiety reduced after the intervention among men and women ($P < 0.001$). In addition, the severity of anxiety among women in the control group increased after placebo intervention ($P < 0.05$). Moreover, the effect of *C. aurantium* aroma was further observed in women than men; however, there was no statistically significant difference ($P > 0.05$).

Conclusions: The results of this study indicated that inhaling the *C. aurantium* aroma reduces anxiety in patients. Aromatherapy also has the potential to be replaced by chemical drugs due to fewer side effects and risks, and cost-effectiveness.

Keywords: Aromatherapy, Anxiety, Operative, Anxiety, Operative; Anxiety; *Citrus aurantium* Essential Oil

1. Background

Surgery is a very stressful condition for patients. Preoperative anxiety is present in 80% of patients (1). Numerous studies have shown that women suffer from anxiety disorders more than men, which may be related to their genetics and hormones (2). The most common causes of anxiety in surgical patients are postoperative complications and fear of medical team mistakes, not regaining consciousness, death, and unknown reasons (3). Preoperative anxiety increases the use of sedatives, pain, nausea, and vomiting after surgery, and risk of infection, and prolongs the patient recovery period (4). On the other hand, one of the effective factors on anxiety is age. Adults become anxious

due to the cost of treatment and length of hospital stay. Hospitalization history and experiences gained from family and friends also affect patients' anxiety levels (5).

Common strategies for the management of preoperative anxiety are pharmacological and non-pharmacological interventions. Effective non-pharmacological interventions in reducing anxiety include acupuncture, massage therapy, music therapy, and aromatherapy (4, 6). Recently, aromatherapy has attracted more attention as a non-pharmacological method to reduce anxiety in patients. The benefits of aromatherapy are low risk, good efficiency, and few side effects (7). This method is accepted in many countries as one of the duties of nurses (8). In aromatherapy, aromatic essential oils

stimulate the olfactory system and release neurotransmitters, cause individuals to calm and reduce anxiety (9).

Aromatherapy reduces anxiety by lowering cortisol levels and increasing serotonin levels. There is evidence that inhaling essential oils releases endorphins and reduces anxiety (10). One of the scents used to reduce anxiety is the scent of *Citrus aurantium*. Citrus essential oils have a stimulating effect on the central nervous system. These essential oils have antispasmodic and anti-inflammatory properties and reduce blood pressure. The flavonoids in *C. aurantium* act as benzodiazepine receptors and reduce anxiety levels. Citrus aroma has the same effect as diazepam and can be substituted (7).

Anxiety disorders are more common in women than men, and these anxiety disorders in women often become more severe during hormonal fluctuations, including puberty period, before menstruation, in pregnancy, or after childbirth and menopause (11); therefore, the incidence of anxiety can also depend on gender (2). The results of a study conducted on the effect of aromatherapy in reducing anxiety in pregnant women showed that aromatherapy decreases anxiety in women (12). Another study also reported the anti-anxiety effects of *C. aurantium* aroma in the first stage of labor (13).

A clinical study showed that aromatherapy with *C. aurantium* reduces anxiety in patients with acute coronary syndrome (14). A clinical trial also showed that aromatherapy with Citrus aurantium is effective in reducing anxiety in patients undergoing coronary angiography (7).

Anxiety disorders are more common in women than men, and these anxiety disorders in women often become more severe during hormonal fluctuations, including puberty period, before menstruation, in pregnancy, or after childbirth and menopause. Numerous studies have shown that women suffer from anxiety disorders more than men, which may be related to their genetics and hormones.

Aromatherapy reduces anxiety by lowering cortisol levels and increasing serotonin levels. There is evidence that inhaling essential oils releases endorphins and reduces anxiety. The results of a study conducted on the effect of aromatherapy in reducing anxiety in pregnant women showed that aromatherapy decreases anxiety in women. Another study also reported the anti-anxiety effects of *C. aurantium* aroma in the first stage of labor.

2. Objectives

The current study aimed to investigate the sedative effects of *C. aurantium* aroma on anxiety in men and women candidates for surgery.

3. Methods

3.1. Trial Design and Participants

This prospective single-blind randomized controlled clinical trial controlled by placebo was performed in hospitals affiliated to Shiraz University of Medical Sciences, Shiraz, Iran, in 2018. The candidates for general surgery were randomly divided into intervention and control groups using a simple lottery method. The total number of participants was 60, with 15 men and 15 women in each group of 30 subjects (i.e., aromatherapy and control groups).

3.2. Ethical Considerations

This study was registered with the ethics code of IR.SUMS.REC.1395.169 in the Ethics Committee of Shiraz University of Medical Sciences and with the code of IRCT2017031433083N1 in the list of clinical trials in Iran. The purpose of the study was explained to the patients. After the approval of the Ethics Committee, written informed consent was obtained from all the participants.

3.3. Sample Size Calculation

The total number of participants in this study was 60. Based on the sample size formula comparing the two means, 30 subjects were determined in each group. The error was considered to be 0.05 with a 95% confidence interval. It should be noted that all the patients participated up to the end of the study (Figure 1).

3.4. Inclusion and Exclusion Criteria

The inclusion criteria were conscious men and women of 40 years and older who were candidates for general surgery, not taken herbal and anti-anxiety drugs before surgery, not addicted to drugs, without employment in the health care system, with no treatment, with no hearing, vision, or smell problems, with no acute illness, and with scores of at least 20 (mild anxiety) in the Spielberger State-Trait Anxiety Inventory (STAI) questionnaire. The exclusion criteria were acute pain at the time of completing the questionnaire, willingness to withdraw from the study at any stage of the intervention, severe anxiety, and emergency surgery.

3.5. Intervention

The patients completed the STAI questionnaire before the intervention. The intervention was performed 60 min before the start of surgery. Two drops of *C. aurantium* essential oil with 100% purity were poured on a napkin using a standard dropper, and the patient inhaled it for 20 min at a distance of 20 cm. The same method was performed in the control group; however, distilled water was used instead

Figure 1. CONSORT flow diagram of study participants

of aroma. The data on overt and covert anxiety were collected using the STAI questionnaire 20 min after the intervention. Evaluations before and after the intervention in all the participants were performed by one individual. In addition, the *C. aurantium* essential oil used in the present study was produced by Barij Essence Pharmaceutical Company in Iran. This essential oil was purified and analyzed using Cooper J.C. chromatography.

3.6. Demographic Information Questionnaire

The patients' demographic information questionnaire, including age, gender, marriage, level of education, and underlying diseases, was completed by the participants before the intervention.

3.7. Spielberger State-Trait Anxiety Inventory

The STAI is often used to assess state and trait anxiety. The STAI scale is valid and reliable, with 40 items, namely 1 - 20 and 21 - 40 related to state and trait anxiety, respectively. Item scoring related to state anxiety is based on a 4-point Likert scale, including (1) not at all, (2) somewhat, (3) moderately so, and (4) very much so. Trait anxiety items are also scored based on a 4-point Likert scale, including (1) almost never, (2) sometimes, (3) often, and (4) almost always (15). Minimum and maximum scores of 20 and 80 are considered for both state anxiety and trait anxiety. The scores obtained from the questionnaire are divided into low (20 - 39), moderate (59 - 40), and high (80 - 60) anxiety. The reliability of this questionnaire, using Cronbach's alpha coefficient, was reported as 93%. The STAI, used nationally and

internationally, has good validity and reliability (4, 16).

3.8. Safety Evaluation

The patients were asked to report side effects at each stage of the study.

3.9. Statistical Analysis

The data obtained from the patients' questionnaire were analyzed by SPSS software (version 16). Descriptive statistics were used to obtain the mean, standard deviation, and frequency distribution tables. Moreover, analytical statistical methods were utilized to perform independent *t*-test, paired *t*-test, and chi-square test. Due to the fact that the normal distribution of the data was obtained using the Kolmogorov-Smirnov test, parametric methods were used for analysis. In addition, a *P*-value of less than 0.05 was considered statistically significant for all the tests.

4. Results

In this study, 60 patients who were candidates for surgery were divided into two groups of intervention (*n* = 30) and control (*n* = 30). Each group consisted of 15 men and 15 women. There was no statistically significant difference between the groups in terms of demographic information (*P* > 0.05; Table 1).

After aromatherapy, more subjects among men reached the level of relatively mild anxiety. Furthermore, most subjects among women reached the level of relatively mild anxiety (Tables 2 and 3). In men and women, the mean score of state anxiety in the aromatherapy group with Citrus aurantium was significantly lower than that reported after the intervention (*P* = 0.001). On the other hand, in women, the mean score of state anxiety in the control group was significantly higher than that reported after placebo intervention (*P* = 0.02; Table 4; Figure 2). In male and female participants, the mean score of trait anxiety in the aromatherapy group with *C. aurantium* was significantly lower than that reported after the intervention (*P* = 0.001; Table 5; Figure 3). The comparison of the mean changes of state and trait anxiety before and after the intervention in the two groups in men and women according to the significance level of the test, which was less than the assumed error in the study, showed significant differences between the groups (Table 6).

5. Discussion

The current study aimed to investigate the anti-anxiety effects of *C. aurantium* aromatherapy on men and women

candidates for surgery. Since one of the factors affecting anxiety is age (5), in order to limit the interfering factor, the patients aged 40 years and older were included in the present study. According to the present study, it can be concluded that inhaling two drops of *C. aurantium* essential oil for 20 min with 100% purity at a distance of 20 cm from the nose was effective in the reduction of anxiety in male and female patients. According to a study conducted on more than 15,000 female patients undergoing non-obstetric surgery, anxiety is often listed as the worst aspect of the preoperative period (17).

Because anxiety is a mental process, psychological factors, such as treatment outcomes, patient characteristics, financial and moral support of the family, and health insurance, also affect the anxiety level of patients (7). In the current study, all the patients had insurance coverage, and none of them had a previous history of surgery because failure or dissatisfaction with previous surgery may increase patient anxiety (18). The findings of a study showed that aromatherapy with *C. aurantium* essential oil reduces anxiety in patients before surgery (4). The results of a study conducted by Pimenta et al. aiming to determine the effect of aromatherapy with *C. aurantium* on anxiety in patients with leukemia before bone marrow removal showed that anxiety levels and physiological parameters significantly improved (19).

In addition, based on a study performed by Moradi et al. on 80 patients, it was concluded that aromatherapy with Citrus aurantium 60 min before coronary angiography for 10 - 15 min reduces the symptoms of anxiety in patients (7). Regarding the improvement of anxiety symptoms, the results of all the aforementioned studies are consistent with the results of the present study. The findings of a review of clinical trials conducted by Manucci et al. showed that the oral administration or inhalation of *C. aurantium* could have beneficial effects on anxiety reduction (20). The main component of *C. aurantium* essential oil is limonene, with anti-anxiety and sedative effects, affecting the activity of the central nervous system (21).

In one study, orange essential oil, with similar compounds to *C. aurantium* essential oil, reduced anxiety in children referring to the dentist (22). In another study, anxiety before coronary artery bypass grafting was reported to be higher in men than that in men (23). Moreover, in a study carried out by Altan et al., anxiety during dental surgery was more frequently reported in women than that in men; however, there was no statistically significant difference (24). Furthermore, in the current study, anxiety before the intervention was higher in women than that in men, but with no statistical significance. In addition, after the intervention with *C. aurantium* essential oil, women's anxiety decreased less than men's anxiety; nevertheless, it

Table 1. Patients' Demographic Data ^a

Variables	Intervention Group	Control Group	P-Value
Total			
Total number of participants	30 (50)	30 (50)	
Gender			
			1.00 ^a
Male	15 (50)	15 (50)	
Female	15 (50)	15 (50)	
Educational level			
			0.920 ^a
Illiteracy	18 (60)	18 (60)	
Under diploma	6 (20)	5 (16.7)	
Above diploma	6 (20)	7 (23.3)	
Marital status			
			1.00 ^a
Married	20 (66.7)	20 (66.7)	
Single	10 (33.3)	10 (33.3)	
Hospitalization			
			0.606 ^a
Yes	16 (53.3)	14 (46.7)	
No	14 (46.7)	16 (53.3)	
Occupancy			
			0.714 ^a
Unemployed	4 (13.3)	5 (16.7)	
Employed	14 (46.7)	16 (53.3)	
Housewife	12 (40)	9 (30)	
Age; mean $\pm$ SD	51.93 $\pm$ 7.26	51.47 $\pm$ 5.87	0.906 ^b

^a Values are expressed as No. (%) unless otherwise indicated.^b Chi-square test.^c Kruskal-Wallis test.**Table 2.** Frequency Distribution of State Anxiety in Two Groups Before and After Intervention ^a

Gender/Anxiety	Control		Citrus aurantium	
	Before Intervention	After Intervention	Before Intervention	After Intervention
Male				
Mild	4 (26.7)	3 (20)		3 (20)
Relatively mild	9 (60)	9 (60)	12 (80)	12 (80)
Relatively intense	2 (13.3)	3 (20)	3 (20)	
Female				
Mild	1 (6.7)			3 (20)
Relatively mild	11 (73.3)	13 (86.7)	12 (80)	11 (73.3)
Relatively intense	3 (20)	2 (13.3)	3 (20)	1 (6.7)
Total				
Mild	5 (16.7)	3 (10)		6 (20)
Relatively mild	20 (66.7)	22 (73.3)	24 (80)	23 (76.7)
Relatively intense	5 (16.7)	5 (16.7)	6 (20)	1 (3.3)

^a Values are expressed as No. (%).

Table 3. Frequency Distribution of Trait Anxiety in Two Groups Before and After Intervention

Gender/Anxiety	Control		<i>Citrus aurantium</i>	
	Before Intervention	After Intervention	After Intervention	Before Intervention
Male				
Mild	3 (20)	3 (20)		3 (20)
Relatively mild	9 (60)	9 (60)	10 (66.7)	12 (80)
Relatively intense	3 (20)	3 (20)	5 (33.3)	
Female				
Mild	1 (6.7)			3 (20)
Relatively mild	12 (80)	13 (86.7)	11 (73.3)	11 (73.3)
Relatively intense	2 (13.3)	2 (13.3)	4 (26.7)	1 (6.7)
Total				
Mild	4 (13.3)	3 (10)		6 (20)
Relatively mild	21 (70)	22 (73.3)	21 (70)	23 (76.7)
Relatively intense	5 (16.7)	5 (16.7)	9 (30)	1 (3.3)

Table 4. Analytical Statistics of State Anxiety in Patients Undergoing Surgery in Two Groups and Comparison of Means at Two Times

Gender/Group/Time	Mean (Standard Deviation)	t-test	Df	P-Value ^a
Male				
Control		-0.50	14	0.628
Before intervention	37.53 (11)			
After intervention	37.87 (9.85)			
<i>Citrus aurantium</i>		11.68	14	0.001
Before intervention	43.53 (8.16)			
After intervention	30.33 (5.73)			
Female				
Control		-2.62	14	0.02
Before intervention	41.87 (8.77)			
After intervention	43.53 (8.98)			
<i>Citrus aurantium</i>		15.33	14	0.001
Before intervention	42.80 (7.88)			
After intervention	30.00 (5.66)			
Total				
Control		-2.12	29	0.043
Before intervention	39.70 (10.02)			
After intervention	40.70 (9.7)			
<i>Citrus aurantium</i>		18.80	29	0.001
Before intervention	43.17 (7.89)			
After intervention	30.17 (5.6)			

^a Paired t-test.

Figure 2. Mean of state anxiety in study groups

Figure 3. Mean of trait anxiety in study groups

Table 5. Analytical Statistics of Trait Anxiety in Patients Undergoing Surgery in Two Groups and Comparison of Means at Two Times

Gender/Group/Time	Mean (Standard Deviation)	t-test	Df	P-Value ^a
Male				
Control		-1.07	14	0.305
Before intervention	38.20 (9.86)			
After intervention	38.60 (9.66)			
<i>Citrus aurantium</i>		20.07	14	0.001
Before intervention	46.33 (7.72)			
After intervention	36.07 (7.29)			
Female				
Control		-1.50	14	0.156
Before intervention	41.87 (8.54)			
After intervention	42.47 (8.11)			
<i>Citrus aurantium</i>		14.81	14	0.001
Before intervention	44.33 (7.99)			
After intervention	33.73 (7.52)			
Total		-1.85	29	0.074
Control				
Before intervention	40.03 (9.25)			
After intervention	40.53 (8.98)			
<i>Citrus aurantium</i>		24.08	29	0.001
Before intervention	45.33 (7.79)			
After intervention	34.90 (7.37)			

^a Paired t-test.

was not statistically significant.

In general, aromatherapy with *C. aurantium* reduced state and trait anxiety significantly in men and women, compared to anxiety before the intervention. In the control group, after the intervention with placebo (i.e., distilled water), state anxiety increased in men; nonetheless, it was not significant. Moreover, trait anxiety remained unchanged. State and trait anxiety also increased significantly in female patients. Therefore, it can be concluded that the placebo increased anxiety in women. However, the results of a study performed by Franco et al. showed that a placebo reduced patients' anxiety, the reason for which could be related to paying too much attention to patients (25). In the present study, no aromatherapy side effects were observed in the patients, which was similar to the results of other studies (4, 21).

According to the results of the current study, the *C. aurantium* essential oil can be used to reduce anxiety in patients waiting for surgery. The strength of this study was the comparison of the aromatherapy effectiveness in men

and women for the first time. The limitations of this study were the lack of odor inhalation by the control group because only the patients in the aromatherapy group inhaled the odor; nonetheless, the control group inhaled distilled water, which may affect the evaluation of the results. On the other hand, the sample size of this study was small, and the study was limited to one city. Therefore, it is recommended to perform studies with a larger sample size in other regions and use different doses and types of fragrances.

5.1. Conclusion

The results of this study indicated that inhaling the *C. aurantium* aroma significantly reduced state and trait anxiety in men and women compared to anxiety before the intervention. Aromatherapy has the potential to be replaced by chemical drugs due to fewer side effects and risks, and cost-effectiveness. Therefore, it is possible to use preoperative aromatherapy performed by a nurse, benefited from a health care program.

Table 6. Comparison of Mean and Standard Deviation Changes of State and Trait Anxiety of Patients Undergoing Surgery in Two groups Before and After Intervention

Variables/Gender/Group	Mean (Standard Deviation)	T	Df	P-Value ^a
State anxiety				
Male		10.283	28	0.001
Control	0.33 (2.61)			
<i>Citrus aurantium</i>	-13.20 (4.38)			
Female		13.772	28	0.001
Control	1.67 (2.47)			
<i>Citrus aurantium</i>	-12.80 (3.23)			
Total		16.719	58	0.001
Control	1.00 (2.59)			
<i>Citrus aurantium</i>	-13.00 (3.79)			
Trait anxiety				
Male		16.812	28	0.001
Control	0.40 (1.45)			
<i>Citrus aurantium</i>	-10.27 (1.98)			
Female		13.659	28	0.001
Control	0.60 (1.55)			
<i>Citrus aurantium</i>	-10.60 (2.77)			
Total		21.411	58	0.001
Control	0.50 (1.48)			
<i>Citrus aurantium</i>	-10.43 (2.37)			

^a Independent t-test.

Footnotes

Authors' Contribution: MA and AE prepared the first draft of the manuscript. MA and JE made critical revisions to the paper and responded to reviewers. ID and KM helped with data analysis.

Clinical Trial Registration Code: This study was registered with a clinical trial registration code (IRCT2017031433083N1).

Conflict of Interests: The authors declare that there is no conflict of interest.

Ethical Approval: The present study was approved by the Ethics Committee of Shiraz University of Medical Sciences (ethics code: IR.SUMS.REC.1395.169).

Funding/Support: This study was financially supported by Shiraz University of Medical Sciences.

Informed Consent: Written informed consent was obtained from all the participants.

References

1. Tulloch I, Rubin JS. Assessment and management of preoperative anxiety. *J Voice*. 2019;**33**(5):691-6. doi: [10.1016/j.jvoice.2018.02.008](https://doi.org/10.1016/j.jvoice.2018.02.008). [PubMed: [29753446](https://pubmed.ncbi.nlm.nih.gov/29753446/)].
2. Scariot R, Tomaz CO, Calixto RD, Gerber JT, Pivetta Petinati MF, Cavalcante RC, et al. Association between gender, estrogen receptors genes and anxiety levels in patients undergoing orthognathic surgery. *J Craniomaxillofac Surg*. 2019;**47**(8):1300-5. doi: [10.1016/j.jcms.2019.05.003](https://doi.org/10.1016/j.jcms.2019.05.003). [PubMed: [31331849](https://pubmed.ncbi.nlm.nih.gov/31331849/)].
3. Pasyar N, Rambod M, Araghi F. The effect of bergamot orange essence on anxiety, salivary cortisol, and alpha amylase in patients prior to laparoscopic cholecystectomy: A controlled trial study. *Complement Ther Clin Pract*. 2020;**39**:101153. doi: [10.1016/j.ctcp.2020.101153](https://doi.org/10.1016/j.ctcp.2020.101153). [PubMed: [32379683](https://pubmed.ncbi.nlm.nih.gov/32379683/)].
4. Darvishi I, Khazaei Z, Hosseinkhani A, Ebrahimi A, Eslami J. The effect of aromatherapy using Lavender (*Lavandula angustifolia* Miller) and *Citrus aurantium* L. extracts to treat anxiety of patients undergoing laparoscopic cholecystectomy: A randomized clinical trial in Iran. *Biomed Res Ther*. 2018;**5**(3):2096-110. doi: [10.15419/bmrat.v5i3.423](https://doi.org/10.15419/bmrat.v5i3.423).
5. Fayazi S, Babashahi M, Rezaei M. The effect of inhalation aromatherapy on anxiety level of the patients in preoperative period. *Iran J Nurs Midwifery Res*. 2011;**16**(4):278-83. [PubMed: [23449862](https://pubmed.ncbi.nlm.nih.gov/23449862/)]. [PubMed Central: [PMC3583096](https://pubmed.ncbi.nlm.nih.gov/PMC3583096/)].
6. Akbarzadeh M, Eslami J, Darvishi I, Ebrahimi A. Effect of acupuncture in Pericardium 6 acupoint on nausea and vomiting after general surgery. *Nursing and Midwifery Studies*. 2019;**8**(3):143. doi: [10.4103/nms.nms_4_18](https://doi.org/10.4103/nms.nms_4_18).

7. Moradi K, Ashtarian H, Danzima NY, Saeedi H, Bijan B, Akbari F, et al. Essential oil from Citrus aurantium alleviates anxiety of patients undergoing coronary angiography: A single-blind, randomized controlled trial. *Chin J Integr Med*. 2021;**27**(3):177–82. doi: [10.1007/s11655-020-3266-5](#). [PubMed: [32572778](#)].
8. Abdelhakim AM, Hussein AS, Doheim MF, Sayed AK. The effect of inhalation aromatherapy in patients undergoing cardiac surgery: A systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med*. 2020;**48**:102256. doi: [10.1016/j.ctim.2019.102256](#). [PubMed: [31987220](#)].
9. Zamanifar S, Bagheri-Saveh MI, Nezakati A, Mohammadi R, Seidi J. The effect of music therapy and aromatherapy with chamomile-lavender essential oil on the anxiety of clinical nurses: A randomized and double-blind clinical trial. *J Med Life*. 2020;**13**(1):87–93. doi: [10.25122/jml-2019-0105](#). [PubMed: [32341707](#)]. [PubMed Central: [PMC7175441](#)].
10. Akbari F, Rezaei M, Khatony A. Effect of peppermint essence on the pain and anxiety caused by intravenous catheterization in cardiac patients: A randomized controlled trial. *J Pain Res*. 2019;**12**:2933–9. doi: [10.2147/JPR.S226312](#). [PubMed: [31695482](#)]. [PubMed Central: [PMC6814313](#)].
11. Hantsoo L, Epperson CN. Anxiety disorders among women: A female lifespan approach. *Focus (Am Psychiatr Publ)*. 2017;**15**(2):162–72. doi: [10.1176/appi.focus.20160042](#). [PubMed: [28966563](#)]. [PubMed Central: [PMC5613977](#)].
12. Igarashi T. Physical and psychologic effects of aromatherapy inhalation on pregnant women: A randomized controlled trial. *J Altern Complement Med*. 2013;**19**(10):805–10. doi: [10.1089/acm.2012.0103](#). [PubMed: [23410527](#)]. [PubMed Central: [PMC3804257](#)].
13. Namazi M, Amir Ali Akbari S, Mojab F, Talebi A, Alavi Majd H, Janesari S. Aromatherapy with citrus aurantium oil and anxiety during the first stage of labor. *Iran Red Crescent Med J*. 2014;**16**(6):e18371. doi: [10.5812/ircmj.18371](#). [PubMed: [25068058](#)]. [PubMed Central: [PMC4102991](#)].
14. Moslemi F, Alijaniha F, Naseri M, Kazemnejad A, Charkhkar M, Heidari MR. Citrus aurantium aroma for anxiety in patients with acute coronary syndrome: A double-blind placebo-controlled trial. *J Altern Complement Med*. 2019;**25**(8):833–9. doi: [10.1089/acm.2019.0061](#). [PubMed: [31211612](#)].
15. Spielberger CD. State-trait anxiety inventory. In: Weiner IB, Craighead WE, editors. *The corsini encyclopedia of psychology*. New Jersey, USA: John Wiley & Sons; 2010. doi: [10.1002/9780470479216.corpsy0943](#).
16. Bech P, Gormsen L, Loldrup D, Lunde M. The clinical effect of clomipramine in chronic idiopathic pain disorder revisited using the Spielberger State Anxiety Symptom Scale (SSASS) as outcome scale. *J Affect Disord*. 2009;**119**(1-3):43–51. doi: [10.1016/j.jad.2009.03.007](#). [PubMed: [19349078](#)].
17. Walker EMK, Bell M, Cook TM, Grocott MPW, Moonesinghe SR; Central Snap-Organisation, et al. Patient reported outcome of adult perioperative anaesthesia in the United Kingdom: A cross-sectional observational study. *Br J Anaesth*. 2016;**117**(6):758–66. doi: [10.1093/bja/aew381](#). [PubMed: [27956674](#)].
18. Kritsidima M, Newton T, Asimakopoulou K. The effects of lavender scent on dental patient anxiety levels: A cluster randomised-controlled trial. *Community Dent Oral Epidemiol*. 2010;**38**(1):83–7. doi: [10.1111/j.1600-0528.2009.00511.x](#). [PubMed: [19968674](#)].
19. Pimenta FC, Alves MF, Pimenta MB, Melo SA, de Almeida AA, Leite JR, et al. Anxiolytic effect of Citrus aurantium L. on patients with chronic myeloid leukemia. *Phytother Res*. 2016;**30**(4):613–7. doi: [10.1002/ptr.5566](#). [PubMed: [26787366](#)].
20. Mannucci C, Calapai F, Cardia L, Inferrera G, D'Arena G, Di Pietro M, et al. Clinical pharmacology of Citrus aurantium and Citrus sinensis for the treatment of anxiety. *Evid Based Complement Alternat Med*. 2018;**2018**:3624094. doi: [10.1155/2018/3624094](#). [PubMed: [30622597](#)]. [PubMed Central: [PMC6304613](#)].
21. Heydari N, Abootalebi M, Jamalimoghadam N, Kasraeian M, Emamghoreishi M, Akbarzadeh M. Investigation of the effect of aromatherapy with Citrus aurantium blossom essential oil on premenstrual syndrome in university students: A clinical trial study. *Complement Ther Clin Pract*. 2018;**32**:1–5. doi: [10.1016/j.ctcp.2018.04.006](#). [PubMed: [30057033](#)].
22. Jafarzadeh M, Arman S, Pour FF. Effect of aromatherapy with orange essential oil on salivary cortisol and pulse rate in children during dental treatment: A randomized controlled clinical trial. *Adv Biomed Res*. 2013;**2**:10. doi: [10.4103/2277-9175.107968](#). [PubMed: [23930255](#)]. [PubMed Central: [PMC3732892](#)].
23. Geulayov G, Novikov I, Dankner D, Dankner R. Symptoms of depression and anxiety and 11-year all-cause mortality in men and women undergoing coronary artery bypass graft (CABG) surgery. *J Psychosom Res*. 2018;**105**:106–14. doi: [10.1016/j.jpsychores.2017.11.017](#). [PubMed: [29332626](#)].
24. Altan A, Akkoc S, Erdil A, Colak S, Demir O, Altan H. Effects of pain catastrophizing and anxiety on analgesic use after surgical removal of impacted mandibular third molars. *J Dent Anesth Pain Med*. 2019;**19**(6):379–88. doi: [10.17245/jdapm.2019.19.6.379](#). [PubMed: [31942453](#)]. [PubMed Central: [PMC6946835](#)].
25. Franco L, Blanck TJ, Dugan K, Kline R, Shanmugam G, Galotti A, et al. Both lavender fleur oil and unscented oil aromatherapy reduce preoperative anxiety in breast surgery patients: A randomized trial. *J Clin Anesth*. 2016;**33**:243–9. doi: [10.1016/j.jclinane.2016.02.032](#). [PubMed: [27555173](#)].